Pythagorean inequalities

In this lesson we'll look at different types of triangles and how to use Pythagorean inequalities to determine what kind of triangle we have based on their angle measures and side lengths.

Types of triangles by angle sizes

Acute triangle

All of the angles are smaller than $90°$

$m\angle A = 79°$

$m\angle B = 45°$

$m\angle C = 56°$



© Krista King Math

Right triangle

The triangle has exactly on right angle

$m\angle A = 28^\circ$

$m\angle B = 90^\circ$

$m\angle C = 62^\circ$



Obtuse triangle

The triangle has one angle greater than 90°

$m\angle A = 45^\circ$

$m\angle B = 110^\circ$

$m\angle C = 25^\circ$

© Krista King Math



Pythagorean inequalities

There is a relationship between the length of the two shortest sides of a triangle and the length of its longest side. If the triangle is not a right triangle, then the relationship is an inequality. Just like in the Pythagorean Theorem, we call the short sides a and b and the long side c.

Acute triangle

$$a^2 + b^2 > c^2$$

© Krista King Math



Right triangle

$$a^2 + b^2 = c^2$$



Obtuse triangle

$$a^2 + b^2 < c^2$$

© Krista King Math



Angles and sides in triangles

One thing to remember about triangles is that the smallest angle is always opposite the shortest side and the biggest angle is always opposite the longest side. Sometimes this can help you when you think through Pythagorean inequality problems.

Let’s start by working through an example.

Example

Classify the triangle with sides 10 in, 5 in, and 9 in as acute, obtuse or right.

Use Pythagorean inequalities to classify the triangles. The two shortest sides are a and b and the longest side is c, so we can say

$$a = 5 \text{ in}$$

$$b = 9 \text{ in}$$

© Krista King Math

$c = 10$ in

Let's see how $a^2 + b^2$ compares with c^2.

$$a^2 + b^2 \stackrel{?}{} c^2$$

$$5^2 + 9^2 \stackrel{?}{} 10^2$$

$$25 + 81 \stackrel{?}{} 100$$

$$106 \stackrel{?}{} 100$$

$$106 > 100$$

Because $a^2 + b^2 > c^2$, this is an acute triangle.

Let's do one more like the first one.

Example

Classify the triangle with sides 9 in, 7 in, and 12 in as acute, obtuse or right.

Use Pythagorean inequalities to classify the triangles. The two shortest sides are a and b and the longest side is c, so we can say

$a = 7$ in

$b = 9$ in

$c = 12$ in

© Krista King Math

Let's see how $a^2 + b^2$ compares with c^2.

$$a^2 + b^2 \stackrel{?}{} c^2$$

$$7^2 + 9^2 \stackrel{?}{} 12^2$$

$$49 + 81 \stackrel{?}{} 144$$

$$130 \stackrel{?}{} 144$$

$$130 < 144$$

Because $a^2 + b^2 < c^2$, this is an obtuse triangle.

Let's try one with a bit more reasoning involved.

Example

A triangle has sides 13 in and 12 in. If the remaining side is the longest side, what is the smallest integer value it can take that would still keep the triangle obtuse?

For a triangle to be obtuse, its sides need to satisfy the inequality $a^2 + b^2 < c^2$. We want to find the largest perfect square that's bigger than $a^2 + b^2$.

We can set $a = 13$ in and $b = 12$ in, so we have

$$13^2 + 12^2$$

© Krista King Math

$$169 + 144$$

$$313$$

We want 313 to be less than c^2, so let's take the square root of 313 and round up to the next integer.

$$\sqrt{313} \approx 17.691$$

This value rounds up to 18, so $c^2 = 18^2 = 324$, and the smallest integer side that makes the triangle obtuse is 18 in.

© Krista King Math